AlphaLISA™ *SureFire*® Biotin Free


# Human PD-L1 Total Detection Kit

**Product number:** ASBF-TPDL1-A500, ASBF-TPDL1-A10K, ASBF-TPDL1-A50K, ASBF-TPDL1-A-HV


**Kit specificity:**
This assay kit contains antibodies which recognize distinct epitopes on PD-L1. The protein detected by this kit corresponds to UniProt ID Q9NZQ7. PD-L1 is also known as PDCD1 ligand 1, Programmed death ligand 1, B7 homolog 1 and CD274. These antibodies recognize PD-L1 of human origin. Other species should be tested on a case-by-case basis.

**Control lysate information:**
Positive Control Lysate: Prepared from A549 cells, cultured to confluence in T175 flasks in 10% FBS containing medium. Cells were treated with 100 ng/mL IFNγ for 48 hours followed by 10 µM ARB-272572 for 15 minutes. Cells were lysed with 4 mL of Lysis Buffer.


**Representative data:**
Data obtained with a 2-plate, 2-incubation protocol. A549 cells were seeded at 20K cells/well in a 96 well plate and incubated overnight. Cells were treated with increasing concentrations of IFNγ for 48 hours. Cells were lysed with Lysis Buffer and assayed separately for Total PD-L1 and ERK using respective *SureFire* Biotin Free kits. Equivalent to approximately 1,000 cells/datapoint.




Data obtained with a 2-plate, 2-incubation protocol. THP-1 cells were seeded at 100K cells/well in a 96 well plate and were treated with increasing concentrations of IFNγ for 48 hours. Cells were lysed with Lysis Buffer and assayed separately for Total PD-L1 and ERK using respective *SureFire* Biotin Free kits. Equivalent to approximately 4,000 cells/datapoint.




Data obtained with a 2-plate, 2-incubation protocol. THP-1 cells were seeded at 1.25 x $10^6$ cells/flask in T25 flasks and were left untreated or treated with 100 ng/mL IFNγ for 48 hours followed by 10 µM of ARB-272572 for 15 minutes. Cells were lysed with Lysis Buffer and assayed separately for PD-L1 Dimer and Total using respective *SureFire* Biotin Free kits. Equivalent to approximately 50,000 cells/datapoint for Dimer or 12,500 cells/datapoint for Total.



Data obtained from measurement of PD-L1 Total in various cell types lysed with Lysis Buffer. Equivalent to approximately 20,000 cells/datapoint for adherent cells or 15,000 cells/datapoint for suspension cells.




The information provided in this document is for reference purposes only and may not be all-inclusive. Revvity, Inc., its subsidiaries, and/or affiliates (collectively, "Revvity") do not assume liability for the accuracy or completeness of the information contained herein. Users should exercise caution when handling materials as they may present unknown hazards. Revvity shall not be liable for any damages or losses resulting from handling or contact with the product, as Revvity cannot control actual methods, volumes, or conditions of use. Users are responsible for ensuring the product's suitability for their specific application. REVVITY EXPRESSLY DISCLAIMS ALL WARRANTIES, INCLUDING WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, REGARDLESS OF WHETHER ORAL OR WRITTEN, EXPRESS OR IMPLIED, ALLEGEDLY ARISING FROM ANY USAGE OF ANY TRADE OR ANY COURSE OF DEALING, IN CONNECTION WITH THE USE OF INFORMATION CONTAINED HEREIN OR THE PRODUCT ITSELF"


Copyright ©2026, Revvity, Inc. All rights reserved.